



The misuse of the Vuong test for non-nested models to test for zero-inflation



Paul Wilson*

School of Mathematics and Computer Science/Statistical Cybermetrics Research Group, University of Wolverhampton, WV1 1LY, United Kingdom

HIGHLIGHTS

- The “Vuong test For Non-Nested Models” is being misused to test zero-inflation.
- This is due to a misunderstanding of the term “non-nested model”.
- And also the mathematically advanced nature of its pre-requisites.
- The hypotheses of the test are misunderstood.
- Allowing negative zero-inflation parameters is a promising alternative.

ARTICLE INFO

Article history:

Received 30 July 2014

Received in revised form

18 October 2014

Accepted 22 December 2014

Available online 31 December 2014

JEL classification:

C52

Keywords:

Non-nested models

Zero-inflation

Vuong test

ABSTRACT

The Vuong test for non-nested models is being widely used as a test of zero-inflation. We show that such use is erroneous. We see that this stems from a misunderstanding of what is meant by the term “non-nested model” and suggest other approaches for determining zero-inflation

© 2014 Elsevier B.V. All rights reserved.

1. The Vuong test for strictly non-nested models

Zero-inflated models are those based upon mixtures of a zero and a count distribution $f(y; \theta)$:

$$f(y; \theta) = \begin{cases} \gamma + (1 - \gamma)f(0; \theta) & y = 0; \\ (1 - \gamma)f(y; \theta) & y = 1, 2, 3, \dots \end{cases} \quad (1)$$

The “Vuong Test for Non-Nested Models” was introduced by Vuong (1989), as a test for “strictly non-nested models”. In slightly simplified form, it states that under the null hypothesis the two non-nested models F_θ and G_τ fit equally well, i.e. the expected value of their log-likelihood ratio equals zero, and then under H_0 the asymptotic distribution of the log-likelihood ratio statistic, LR , is normal. In particular, (under H_0):

$$\frac{LR_n(\hat{\theta}_n, \hat{\tau}_n)}{\hat{\omega}_n \sqrt{n}} \rightarrow N(0, 1) \quad (2)$$

where ω denotes the variance of LR_n and n the sample size. Vuong (1989) also presents tests for nested and overlapping models, and shows that, given certain conditions, the distributions of their log-likelihood ratios are related to χ^2 distributions. Due to the simplicity of its calculation, the test has become popular among statistical practitioners in various disciplines and is implementable in the R-package *pscl*, (Jackman, 2012); Stata offers the Vuong test as an option in its command for zero-inflated Poisson models.

2. The misuse of the Vuong test

The Vuong test for strictly non-nested models is being widely misused as a test of zero-inflation, even though the original paper of Vuong (1989) does not mention zero-inflation. For example the help page associated with the *vuong* command in *pscl* states: “The Vuong non-nested test is based on a comparison of the predicted probabilities of two models that do not nest. Examples include comparisons of zero-inflated count models with their non-zero-inflated analogs (e.g., zero-inflated Poisson versus ordinary Poisson, or zero-inflated negative-binomial versus ordinary negative-binomial).”

* Tel.: +44 1902 321444.

E-mail address: pauljwilson@wlv.ac.uk.




Fig. 1. Distributions of the log-likelihood ratios of ZIP versus Poisson and strictly non-nested models.

Desmarais and Harden (2013) state that: “researchers commonly use the Vuong test (Vuong, 1989) to determine whether the zero-inflated model fits the data statistically significantly better than count regression with a single equation” and cite ten references to publications that have used the Vuong test for this purpose. That this is an incorrect use of the Vuong test for non-nested models is illustrated by Fig. 1. The top histogram illustrates the observed distribution of the log-likelihood ratios obtained when a one-covariate zero-inflated Poisson (ZIP) model and the corresponding Poisson model are fitted to 100,000 samples of size $n = 100$ under the null hypothesis. Clearly the distribution is non-normal. The bottom histogram is produced using exactly the same software code that produced the top histogram, but here the generating model was Poisson with four uniformly distributed covariates x_1 – x_4 , and the competing models were each Poisson with two covariates, x_1, x_2 and x_3, x_4 respectively, and hence are strictly non-nested according to the definition of Vuong (1989).

Desmarais and Harden (2013) discuss AIC and BIC type adjustments to the distribution of the log-likelihood ratios, and present evidence that this improves the power of the Vuong

test; these adjustments are, for any given comparison of models, constants, and hence only effect the mean of the distribution, not its shape.

2.1. The cause of the confusion

The misuse of the test stems from misunderstanding of what is meant by the terms “non-nested model” and “nested model”. As is the case with many frequently used terms their meanings are approximately understood by many, but precisely understood by few. Clarke (2001) observes that “Defining the concept of ‘non-nested’ precisely is not an easy task. Definitions are often imprecise and uncomplicated or precise and complicated.” This statement applies equally well to nested models. (Indeed one category is not the complement of the other; there exists a third, “inbetween” category of what Vuong (1989) refers to as *overlapping models*.) Simple definitions of nested model include that of Davison (2003): “Two models are said to be nested if one reduces to the other when certain parameters are fixed”. In contrast Vuong (1989) defines a model G_F to be nested in a model F_Θ by: “ G_F is nested in F_Θ if and only if $G_F \subset F_\Theta$.” (“ \subset ” indicating that the parameters of G are a proper subset of those of F .)

The distribution of the log-likelihood ratios of non-nested models being normal is dependent on six assumptions presented elsewhere in Vuong (1989). These refer to various topological and measure theoretic properties which are unlikely to be understood except by those with a strong mathematical background. For example: “For every θ in Θ and for H^0_z -almost all z the conditional distribution $F_{Y|Z}(\cdot|z, \Theta)$ has a Radon–Nikodym density $f(\cdot|z, \Theta)$ relative to ν_Y , which is strictly positive for ν_Y -almost all y ”.

Of particular relevance here is the prerequisite of the Vuong test that “ θ_* is an interior point of Θ ”, i.e. nesting must *not* occur at a boundary of the parameter space of the larger model. Whilst the probability distribution function (see Eq. (1)) of a zero-inflated Poisson distribution with zero-inflation parameter γ , where $0 \leq \gamma \leq 1$, reduces to that of a Poisson distribution when $\gamma = 0$, this value is on the boundary of the permissible values of the zero-inflation parameter as γ may not be negative. This results in non-normality of the sampling distribution of the zero-inflation parameter. As Vuong’s subsequent theoretical development of the distributions of log-likelihood ratios depends upon normality of the sampling distribution of the model parameters, clearly his theory is not applicable.

2.2. Models fitted using link functions

Zero-inflated models are usually fitted using a logit link to model the expected proportion of perfect zeros. Whilst it is true that the logistic function: $\frac{\exp(t)}{1+\exp(t)} \neq 0$ for all $t \in \mathbb{R}$, and hence in some sense this formulation of the ZIP and of Poisson are non-nested, $\lim_{t \rightarrow -\infty} \frac{\exp(t)}{1+\exp(t)} = 0$; thus this formulation of the zero-inflated model fails to meet Vuong’s prerequisite that the parameter space is a compact subset of \mathbb{R}^p , and, similar to the scenario presented in the previous section the sampling distribution of the zero-inflation parameter, and hence the distribution of the log-likelihood ratios, is non-normal. Similar statements hold if probit or complementary log–log links are used. It is worth noting that there is confusion in the literature about models being nested if one reduces to the other if certain parameters are fixed, many authors apparently taking this to mean fixed *at zero*. For example, Desmarais and Harden (2013) state: “the count regression f is not nested in the zero-inflated model, because the model does not reduce to f (the count model) when $\gamma = 0$, in which case the probability of a 0 is inflated by 0.50”, apparently alluding to the fact that the value of the logistic function = 0.5 when $t = 0$.

2.3. The null hypothesis of Vuong's test

Consistent with a test of zero-inflation, the simulated distribution of the Vuong statistic presented in the left diagram of Fig. 1 is derived by resampling from non-zero-inflated data. As stated in Section 1 the null hypothesis of Vuong's test for non-nested models is that the expected value of their log-likelihood ratios equals zero. This implies that under the null hypothesis both models are "equally far away" from the data that is being modelled. If we temporarily ignore the issue of whether zero-inflated models and their non-zero-inflated counterparts are non-nested or otherwise, and consider them to be non-nested, to appropriately simulate the distribution of the log-likelihood ratios it would be necessary to resample from data that was somehow equidistant from zero-inflated and non-zero-inflated data; it is difficult to envisage the nature of such data. More importantly, non-rejection of the null hypothesis of Vuong's test for non-nested models, where the (supposedly) non-nested models are, say, the zero-inflated Poisson and standard Poisson model would mean that there is no evidence to conclude that either model fits the data better than the other, not that there is no evidence to support zero-inflation, and its rejection simply implies that either the zero-inflated Poisson model fits the data better than the Poisson model, or vice-versa, not that zero-inflation is present or absent.

3. Other approaches

Distributional methods: early research by the author indicates that if negative values of the log-likelihood ratio that are very close to zero are considered as zeros, then the distribution of ZIP versus Poisson log-likelihood ratios, where the zero inflation parameter is only modelled by an intercept, is a mixture of a point mass at zero and an χ^2_1 distribution, the weighting of the mixture being dependent on the number of covariates. If the zero-inflation parameter is modelled, a mixture of a zero point mass and some other distribution still occurs, whilst the nature of this other distribution is yet to be determined. Self and Liang (1987) discuss the distribution of log-likelihood ratios under non-

standard conditions in a general setting. Note that when the value of γ is allowed to be both positive or negative fitted values of γ do not "pile up" close to zero, and the distribution of zero-modification parameter is normal and hence a non-zero-inflated model is (strictly) nested in its zero-inflated counterpart, and hence a Vuong test for nested models could be used as a test of zero-inflation/deflation. Dietz and Böhning (2000) proposed a link function that allowed for zero-deflation, and more recently Todem et al. (2012) have proposed a score test that incorporates a link function that allows for both zero-inflation and deflation.

4. Conclusion

This article has shown that the widespread practice of using Vuong's test for non-nested models as a test of zero-inflation is erroneous. The misuse is rooted in a misunderstanding of what is meant by the term "non-nested model". The derivation of the distribution of the log-likelihood ratios of zero-inflated versus non-zero inflated models is not straightforward, and a possible alternative approach is to develop tests based upon zero-modified models where the zero-"inflation" parameter may be negative.

References

- Clarke, K., 2001. Testing nonnested models of international relations: re-evaluating realism. *Amer. J. Polit. Sci.* 45, 724–744.
- Davison, A., 2003. *Statistical Models*. Cambridge University Press.
- Desmarais, B.A., Harden, J.J., 2013. Testing for zero inflation in count models: bias correction for the Vuong test. *Stata J.* 13, 810–835.
- Dietz, D., Böhning, D., 2000. On the estimation of the Poisson parameter in zero-modified Poisson models. *Comput. Statist. Data Anal.* 34, 441–459.
- Jackman, S., 2012. *pscl: Classes and Methods for R Developed in the Political Science Computational Laboratory*, Stanford University. Stanford University. R package version 1.04.4. URL: <http://pscl.stanford.edu/>.
- Self, S.G., Liang, K.Y., 1987. Asymptotic properties of maximum likelihood estimators and likelihood ratio tests under nonstandard conditions. *J. Amer. Statist. Assoc.* 82, 605–610.
- Todem, D., Hsu, W., Kim, K., 2012. On the efficiency of score tests for homogeneity in two-component parametric models for discrete data. *Biometrics* 68, 975–982.
- Vuong, Q., 1989. Likelihood ratio tests for model selection and non-nested hypotheses. *Econometrica* 57, 307–333.